

A unified understanding of the two formulae for the traces of the inverse powers of a positive definite symmetric tridiagonal matrix

Takumi Yamashita ¹

Abstract

For an upper bidiagonal matrix B where all the diagonal and the upper sub-diagonal entries are positive, two subtraction-free formulae for computation of the traces $J_M(B) = \text{Tr}((B^\top B)^{-M}) = \text{Tr}((BB^\top)^{-M})$ ($M = 1, 2, \dots$) have been presented in the two preceding works. A few lower bounds of the minimal singular value of B are obtained from these traces. In this paper, we clarify some properties of these formulae and present a new subtraction-free formula for the traces $J_M(B)$. An interpretation of some quantities in one of the preceding works in terms of matrix theory is given. Some relationships between some quantities in the preceding works are also given. From these relationships, the new subtraction-free formula for the traces $J_M(B)$ is obtained.

1 Introduction

Singular values of a matrix are theoretically important and have wide applications. Their exact values are, needless to say, have a lot of examples of applications. In addition to such examples, estimation of a lower bound of the minimal singular value also has examples of applications. An example is estimation of an upper bound of condition number of a matrix. In a system of simultaneous linear equations or an eigenvalue problem, condition number of the coefficient matrix is used as an indicator for reliability of numerical computation. Condition number $\kappa(A)$ of a matrix A is defined by norms of the matrices A and A^{-1} as $\kappa(A) = \|A\| \cdot \|A^{-1}\|$. When we choose the 2-norm for definition of condition number, this value is expressed by the maximal singular value $\sigma_{\max}(A)$ and the minimal singular value $\sigma_{\min}(A)$ of A as $\kappa(A) = \sigma_{\max}(A)/\sigma_{\min}(A)$. The exact values of the maximal and the minimal singular values of A are usually unknown before solving the problem. Then, an upper bound of condition number is used as an alternative in practice. An upper bound of $\sigma_{\max}(A)$ is given as $\sigma_{\max}(A) \leq \sqrt{\|A\|_1 \cdot \|A\|_\infty}$,

¹28-20 Kojogaoka, Otsu, Shiga 520-0821 Japan, Tel.: +81-77-522-7447, Fax: +81-77-522-7447, e-mail: t-yamashita@kke.biglobe.ne.jp

where $\|A\|_1$ and $\|A\|_\infty$ are the 1-norm and the infinity norm of A , respectively [4]. Lower bounds of the minimal singular value of a matrix have been proposed by several authors. See [2, 4, 5, 7, 9, 13], for example. As another example of applications, a lower bound of the minimal singular value of a bidiagonal matrix can be applied to singular value computing. In standard procedure for singular value computing, the input matrix is firstly transformed into a bidiagonal matrix by orthogonal transformations without changing its singular values. Then, we obtain the singular values by applying some algorithm to this bidiagonal matrix. For example, by the dqds (the differential quotient-difference algorithm with shifts) algorithm [1], the orthogonal qd-algorithm [8], or the mdLVs algorithm (modified discrete Lotka-Volterra algorithm with shift) [3], we can obtain singular values of a bidiagonal matrix. These algorithms are iterative and equip a technique called the shift of origin. This technique accelerates convergence of iteration. In this technique, we are required to give a quantity called a shift at each iteration. A too large shift causes failure of the algorithms. A shift can be determined from a lower bound of the minimal singular value of the bidiagonal matrix. Kimura *et al.* [6] presented a sequence of lower bounds of the minimal singular value $\sigma_{\min}(B)$ of an upper bidiagonal matrix B , where all the diagonal and the upper subdiagonal entries are positive. Let these lower bounds be denoted by $\theta_M(B)$ for $M = 1, 2, \dots$. They are given as

$$\theta_M(B) = (J_M(B))^{-\frac{1}{2M}},$$

where $J_M(B) = \text{Tr}((B^\top B)^{-M})$. Note that it holds $\text{Tr}((B^\top B)^{-M}) = \text{Tr}((BB^\top)^{-M})$. These lower bounds increase monotonically and converges to the minimal singular value $\sigma_{\min}(B)$ as $M \rightarrow \infty$ [6, Theorem 3.1], namely,

$$\begin{aligned} \theta_1(B) &< \theta_2(B) < \dots < \sigma_{\min}(B), \\ \lim_{M \rightarrow \infty} \theta_M(B) &= \sigma_{\min}(B). \end{aligned}$$

Kimura *et al.* [6] presented also a formula for computation of the trace $J_M(B)$ for a fixed positive integer M . This formula gives the diagonal entries of $(B^\top B)^{-M}$ and $(BB^\top)^{-M}$ in a form of recurrence relation. In the case of $M \geq 2$, the recurrence relation includes subtraction in it. Subtraction may occur cancellation error in numerical computation. To exclude this possibility, Yamashita, Kimura and Nakamura [11] derived a “subtraction-free” recurrence formula, which consists of only summation, multiplication and division among positive quantities, for the diagonal entries of $(B^\top B)^{-M}$ and $(BB^\top)^{-M}$ ($M = 1, 2, \dots$) starting from the formula in [6]. On the other hand, Yamashita, Kimura and Yamamoto [12] derived another subtraction-free formula for the traces $J_M(B)$ ($M = 1, 2, \dots$) with an idea which is quite different from that in [11]. This derivation does not aim to obtain the diag-

onal entries of $(B^\top B)^{-M}$ or $(BB^\top)^{-M}$ ($M = 1, 2, \dots$). These formulae in [6], [11] and [12] are briefly reviewed in the next section.

In this paper, we consider the two formulae in the preceding works [11] and [12] and achieve a unified understanding of them. Firstly, we show that some quantities used in the formula in [11] can be interpreted in terms of matrix theory. Secondly, we reveal relationships between some quantities in the two formulae. Lastly, a new subtraction-free formula for the traces $J_M(B)$ ($M = 1, 2, \dots$) is obtained from these relationships.

This paper is organized as follows. In Section 2, we briefly review the formulae in the preceding works [6], [11] and [12]. In Section 3, we achieve the unified understanding of the formulae in [11] and [12] and present the new formula. Section 4 is devoted for concluding remarks.

2 The formulae for the traces in the preceding works

In this section, we briefly review the formulae for the traces $J_M(B)$ ($M = 1, 2, \dots$) in the preceding works [6], [11] and [12]. The review of [6] helps to explain the formula in [11]. Before review, notation of the $N \times N$ upper bidiagonal matrix $B = (B_{i,j})$, where all the diagonal and the upper subdiagonal entries are positive, is necessary. Let the matrix B be denoted by

$$B = \begin{pmatrix} \sqrt{q_1} & \sqrt{e_1} & & & \\ & \sqrt{q_2} & \sqrt{e_2} & & \\ & & \ddots & \ddots & \\ & & & \sqrt{q_{N-1}} & \sqrt{e_{N-1}} \\ & & & & \sqrt{q_N} \end{pmatrix}, \quad (1)$$

where $q_i > 0$ for $i = 1, \dots, N$ and $e_i > 0$ for $i = 1, \dots, N - 1$. When we consider singular values of an upper bidiagonal matrix, we can impose the condition of positivity of the diagonal and the upper subdiagonal entries without loss of generality [1]. The preceding works [6] and [11] present formulae for computation of the diagonal entries of $(B^\top B)^{-M}$ and $(BB^\top)^{-M}$ ($M = 1, 2, \dots$). As the sum of these entries, the trace $J_M(B)$ is obtained. These formulae are reviewed in Sections 2.1 and 2.2. On the other hand, the derivation of the formula in the preceding work [12] does not aim to obtain the diagonal entries of $(B^\top B)^{-M}$ or $(BB^\top)^{-M}$ ($M = 1, 2, \dots$). This formula is reviewed in Section 2.3. Note that we use the convention $\sum_{i=j}^k = 0$ if $j > k$ throughout this paper.

2.1 A review of the formula in the preceding work [6]

In this subsection, we briefly review the formula for the trace $J_M(B)$ for a fixed positive integer M in the preceding work by Kimura *et al.* [6]. This

formula gives the diagonal entries of the matrices $(B^\top B)^{-M}$ and $(BB^\top)^{-M}$ in a form of recurrence relation. As the sum of these diagonal entries, the trace $J_M(B)$ is obtained. For a fixed positive integer M and $m = 0, 1, \dots, M$, let the (i, i) -entry of the matrices $(B^\top B)^{-m}$ and $(BB^\top)^{-m}$ be denoted by $v_i^{(m)}$ and $w_i^{(m)}$, respectively. These diagonal entries are obtained by the following theorem.

Theorem 1

Let M be a fixed positive integer. All the diagonal entries $v_i^{(M)}$ and $w_i^{(M)}$ of inverse matrices $(B^\top B)^{-M}$ and $(BB^\top)^{-M}$, respectively, are obtained through a finite number of arithmetics by using the following simple recurrence relations. The recurrence relations are

$$v_i^{(0)} = 1, \quad i = 1, \dots, N, \quad (2)$$

$$w_i^{(0)} = 1, \quad i = 1, \dots, N, \quad (3)$$

$$v_N^{(p)} = \frac{1}{q_N} w_N^{(p-1)}, \quad (4)$$

$$v_i^{(p)} = \frac{1}{q_i} (e_i v_{i+1}^{(p)} + z_i^{(p-1)} - w_i^{(p-1)}), \quad i = 1, \dots, N-1, \quad (5)$$

$$w_1^{(p)} = \frac{1}{q_1} v_1^{(p-1)}, \quad (6)$$

$$w_i^{(p)} = \frac{1}{q_i} (e_{i-1} w_{i-1}^{(p)} + z_i^{(p-1)} - v_i^{(p-1)}), \quad i = 2, \dots, N, \quad (7)$$

$$z_1^{(q)} = 2v_1^{(q)}, \quad (8)$$

$$z_i^{(q)} = z_{i-1}^{(q)} + 2(v_i^{(q)} - w_{i-1}^{(q)}), \quad i = 2, \dots, N \quad (9)$$

for integers p and q such that $1 \leq p \leq M$ and $0 \leq q \leq M-1$. Instead of equations (8) and (9), the following relations can be used:

$$z_N^{(q)} = 2w_N^{(q)},$$

$$z_i^{(q)} = z_{i+1}^{(q)} + 2(w_i^{(q)} - v_{i+1}^{(q)}), \quad i = 1, \dots, N-1$$

for integers q such that $0 \leq q \leq M-1$.

Substitution of $p = 1$ and $q = 0$ into the recurrence relations in Theorem 1 gives the following remark.

Remark 2

The recurrence relations from (4) through (7) in Theorem 1 for $p = 1$ are

simplified to the recurrence relations

$$v_N^{(1)} = \frac{1}{q_N}, \quad (10)$$

$$v_i^{(1)} = \frac{1}{q_i} (e_i v_{i+1}^{(1)} + 1), \quad i = 1, \dots, N-1, \quad (11)$$

$$w_1^{(1)} = \frac{1}{q_1}, \quad (12)$$

$$w_i^{(1)} = \frac{1}{q_i} (e_{i-1} w_{i-1}^{(1)} + 1), \quad i = 2, \dots, N. \quad (13)$$

Theorem 1 for $M = 1$ is reduced to these recurrence relations.

It is noted that there exist some preceding works on some limited cases in numerical analysis.

Remark 3

A formula related to the equations from (10) to (13) for computing the diagonal entries of the inverse $(BB^\top)^{-1}$ is known. See [1, 8, 10], for example. Another formula for computing the diagonal entries of the inverse $(BB^\top)^{-2}$ is presented by von Matt [8].

2.2 A review of the formula in the preceding work [11]

In this subsection, we briefly review the preceding work by Yamashita, Kimura and Nakamura [11]. They derived a formula for computing the diagonal entries of the inverse powers $(B^\top B)^{-M}$ and $(BB^\top)^{-M}$ ($M = 1, 2, \dots$) in a form of recurrence relations starting from the formula in [6] which is reviewed in the previous subsection. In contrast to the formula in [6] which includes subtraction in it in the cases of $M \geq 2$, the derived formula consists of only addition, multiplication and division among positive quantities. Namely, it is “subtraction-free”. Any possibility of cancellation error in numerical computation is clearly excluded from this property. As the sum of these diagonal entries, the traces $J_M(B)$ ($M = 1, 2, \dots$) are obtained.

For $p = 1, \dots, M$, let the (i, i) -entry of $(B^\top B)^{-p}$ and $(BB^\top)^{-p}$ be denoted by $v_i^{(p)}$ and $w_i^{(p)}$, respectively. Quantities $\check{B}_i$ ($i = 1, \dots, N$), F_i ($i = 1, \dots, N-1$) and $\tilde{F}_i$ ($i = 2, \dots, N$) are introduced as follows.

Definition 4

$$\check{B}_i = \frac{1}{q_i}, \quad i = 1, \dots, N, \quad (14)$$

$$F_i = \frac{e_i}{q_i} = e_i \check{B}_i, \quad i = 1, \dots, N-1, \quad (15)$$

$$\tilde{F}_i = \frac{e_{i-1}}{q_i} = e_{i-1} \check{B}_i, \quad i = 2, \dots, N. \quad (16)$$

Next, quantities $g_i^{(r)}$ and $\tilde{g}_i^{(r)}$ for $i = 1, \dots, N$ and $r = 1, 2, \dots$ are introduced as follows.

Definition 5

The quantities $g_i^{(r)}$ for $i = 1, \dots, N$ and $r = 1, 2, \dots$ are defined as follows.

- For $i = N$ and arbitrary positive integer r , $g_N^{(r)}$ is given as $g_N^{(r)} = 0$.
- For $i = 1, \dots, N-1$ and $r = 1$, $g_i^{(1)}$ is given as $g_i^{(1)} = F_i v_{i+1}^{(1)}$.
- For $i = 1, \dots, N-1$ and $r = 2, 3, \dots$, $g_i^{(r)}$ is given as

$$g_i^{(r)} = F_i g_{i+1}^{(r)} + \check{B}_{i+1} g_i^{(r-1)} + \sum_{k=1}^{r-1} g_{i+1}^{(k)} g_i^{(r-k)}. \quad (17)$$

Definition 6

The quantities $\tilde{g}_i^{(r)}$ for $i = 1, \dots, N$ and $r = 1, 2, \dots$ are defined as follows.

- For $i = 1$ and arbitrary positive integer r , $\tilde{g}_1^{(r)}$ is given as $\tilde{g}_1^{(r)} = 0$.
- For $i = 2, \dots, N$ and $r = 1$, $\tilde{g}_i^{(1)}$ is given as $\tilde{g}_i^{(1)} = \tilde{F}_i w_{i-1}^{(1)}$.
- For $i = 2, \dots, N$ and $r = 2, 3, \dots$, $\tilde{g}_i^{(r)}$ is given as

$$\tilde{g}_i^{(r)} = \tilde{F}_i \tilde{g}_{i-1}^{(r)} + \check{B}_{i-1} \tilde{g}_i^{(r-1)} + \sum_{k=1}^{r-1} \tilde{g}_{i-1}^{(k)} \tilde{g}_i^{(r-k)}. \quad (18)$$

Remark 7

The recurrence relations in Remark 2 can be rewritten with the quantities defined by Definitions from 4 to 6 as follows.

$$v_N^{(1)} = \check{B}_N, \quad (19)$$

$$v_i^{(1)} = F_i v_{i+1}^{(1)} + \check{B}_i = g_i^{(1)} + \check{B}_i, \quad i = 1, \dots, N-1, \quad (20)$$

$$w_1^{(1)} = \check{B}_1, \quad (21)$$

$$w_i^{(1)} = \tilde{F}_i w_{i-1}^{(1)} + \check{B}_i = \tilde{g}_i^{(1)} + \check{B}_i, \quad i = 2, \dots, N. \quad (22)$$

A formula for the traces $J_M(B)$ ($M \geq 2$) is given as follows.

Theorem 8

For $M \geq 2$, the diagonal entries $v_i^{(s)}$ and $w_i^{(s)}$ of $(B^\top B)^{-s}$ and $(BB^\top)^{-s}$, respectively, for $i = 1, \dots, N$ and $s = 2, \dots, M$ are computed by the recurrence relations

$$\begin{aligned} v_N^{(s)} &= \check{B}_N w_N^{(s-1)}, \\ w_1^{(s)} &= \check{B}_1 v_1^{(s-1)}, \\ v_i^{(s)} &= F_i v_{i+1}^{(s)} + \check{B}_i w_i^{(s-1)} + 2 \sum_{k=1}^{s-1} g_i^{(k)} w_i^{(s-k)}, \quad i = 1, \dots, N-1, \\ w_i^{(s)} &= \tilde{F}_i w_{i-1}^{(s)} + \check{B}_i v_i^{(s-1)} + 2 \sum_{k=1}^{s-1} \tilde{g}_i^{(k)} v_i^{(s-k)}, \quad i = 2, \dots, N, \end{aligned}$$

with the recurrence relations from (19) to (22).

2.3 A review of the formula in the preceding work [12]

In this subsection, we briefly review the preceding work by Yamashita, Kimura and Yamamoto [12]. Since the derivation of the formula does not aim to obtain the diagonal entries of $(B^\top B)^{-p}$ or $(BB^\top)^{-p}$ ($p = 1, 2, \dots$), we review the idea for the derivation briefly first. Let us consider the matrix $B^\top B$. It is transformed into

$$A = \begin{pmatrix} q_1 & q_1 e_1 & & \\ 1 & q_2 + e_1 & \ddots & \\ & \ddots & \ddots & q_{N-1} e_{N-1} \\ & & 1 & q_N + e_{N-1} \end{pmatrix}$$

by similarity transformation without changing its eigenvalues. Let I be the $N \times N$ unit matrix. In derivation of the formula from the matrix $B^\top B$, one of the key points is to express the determinant $D(\lambda) = \det(A - \lambda I)$ in two ways. For the eigenvalues $\lambda_1, \dots, \lambda_N$ of $B^\top B$, the determinant is given as $D(\lambda) = \prod_{i=1}^N (\lambda_i - \lambda)$. The matrix $A - \lambda I$ can be decomposed as

$$A - \lambda I = \begin{pmatrix} \hat{q}_1^{(0)} & & & \\ 1 & \hat{q}_2^{(0)} & & \\ & \ddots & \ddots & \\ & & 1 & \hat{q}_N^{(0)} \end{pmatrix} \begin{pmatrix} 1 & \hat{e}_1^{(0)} & & \\ & 1 & \ddots & \\ & & \ddots & \hat{e}_{N-1}^{(0)} \\ & & & 1 \end{pmatrix},$$

where $\hat{q}_i^{(0)}$ for $i = 1, \dots, N$ and $\hat{e}_i^{(0)}$ for $i = 1, \dots, N-1$ are functions of λ . The superscript (0) means that these functions have not been differentiated

yet. The other expression of the determinant is $D(\lambda) = \prod_{i=1}^N \hat{q}_i^{(0)}$. Since the matrices $B^\top B$ and A have the same eigenvalues, it holds

$$\prod_{i=1}^N (\lambda_i - \lambda) = \prod_{i=1}^N \hat{q}_i^{(0)}. \quad (23)$$

Let $\hat{q}_i^{(p)}$ for $i = 1, \dots, N$ and $p = 1, 2, \dots$ be $d^p \hat{q}_i^{(0)} / d\lambda^p$. By differentiating (23) with respect to λ , we have

$$\sum_{i=1}^N \left(-\frac{D(\lambda)}{\lambda_i - \lambda} \right) = \sum_{i=1}^N \left(\frac{\hat{q}_i^{(1)}}{\hat{q}_i^{(0)}} \cdot D(\lambda) \right).$$

Then, it holds

$$\sum_{i=1}^N \frac{1}{\lambda_i - \lambda} = \sum_{i=1}^N \left(-\frac{\hat{q}_i^{(1)}}{\hat{q}_i^{(0)}} \right). \quad (24)$$

Let $\hat{H}_i^{(p)}$ for $i = 1, \dots, N$ and $p = 1, 2, \dots$ be functions of λ defined as

$$\hat{H}_i^{(p)} = \begin{cases} -\frac{\hat{q}_i^{(1)}}{\hat{q}_i^{(0)}}, & p = 1, \\ \frac{d\hat{H}_i^{(p-1)}}{d\lambda}, & p = 2, 3, \dots \end{cases} \quad (25)$$

Then, by differentiating (24) repeatedly, we have

$$(p-1)! \sum_{i=1}^N \frac{1}{(\lambda_i - \lambda)^p} = \sum_{i=1}^N \hat{H}_i^{(p)}, \quad p = 1, 2, \dots \quad (26)$$

Thus, the traces of $(B^\top B)^{-p}$ ($p = 1, 2, \dots$) are obtained by substituting $\lambda = 0$ into (26). The result is

$$\text{Tr}((B^\top B)^{-p}) = \frac{1}{(p-1)!} \sum_{i=1}^N H_i^{(p)}, \quad p = 1, 2, \dots, \quad (27)$$

where $H_i^{(p)}$ is $H_i^{(p)} = \hat{H}_i^{(p)}|_{\lambda=0}$. From this result, a method for computing $H_i^{(p)}$ is required. As such a method, a recurrence relation with a help of auxiliary quantities $h_i^{(p)}$ for $i = 1, \dots, N$ and $p = 1, 2, \dots$ is given. The auxiliary quantities $h_i^{(p)}$ is defined as $h_i^{(p)} = \hat{h}_i^{(p)}|_{\lambda=0}$, where $\hat{h}_i^{(p)}$ are functions of λ defined as $\hat{h}_i^{(p)} = -\hat{q}_i^{(p)} / \hat{q}_i^{(0)}$. The recurrence relation is presented as the following theorem.

Theorem 9

Let B be an upper bidiagonal matrix defined in (1).

Let us introduce constants $\tilde{F}_i$ for $i = 2, \dots, N$ defined as

$$\tilde{F}_i = \frac{e_{i-1}}{q_i}, \quad i = 2, \dots, N.$$

For $i = 1, \dots, N$ and $p = 1, 2, \dots$, let $h_i^{(p)}$ be constants which satisfy the following recurrence relation

$$h_1^{(1)} = \frac{1}{q_1}, \quad (28)$$

$$h_i^{(1)} = \tilde{F}_i h_{i-1}^{(1)} + \frac{1}{q_i}, \quad i = 2, \dots, N, \quad (29)$$

$$h_1^{(p)} = 0, \quad p = 2, 3, \dots, \quad (30)$$

$$h_i^{(p)} = \tilde{F}_i \left(h_{i-1}^{(p)} + p h_{i-1}^{(1)} h_{i-1}^{(p-1)} \right) + \sum_{k=1}^{p-2} p C_k h_{i-1}^{(k)} h_i^{(p-k)}, \quad i = 2, \dots, N, \quad p = 2, 3, \dots \quad (31)$$

For $i = 1, \dots, N$ and $p = 1, 2, \dots$, let $H_i^{(p)}$ be constants which satisfy the following recurrence relation

$$H_i^{(1)} = h_i^{(1)}, \quad i = 1, \dots, N, \\ H_i^{(p)} = h_i^{(p)} + \sum_{k=1}^{p-1} C_k h_i^{(k)} H_i^{(p-k)}, \quad i = 1, \dots, N, \quad p = 2, 3, \dots \quad (32)$$

The traces $\text{Tr}((B^\top B)^{-p})$ for $p = 1, 2, \dots$ are computed by

$$\text{Tr}((B^\top B)^{-p}) = \frac{1}{(p-1)!} \sum_{i=1}^N H_i^{(p)}.$$

Remark 10

The recurrence relation of $h_i^{(1)}$ ($i = 1, \dots, N$) in Theorem 9 is equivalent to the recurrence relation for the diagonal entries of $(BB^\top)^{-1}$ shown in Remark 2 (Originally, Remark 4.6 in [6]). Then, the constants $h_i^{(1)}$ and $H_i^{(1)}$ are the (i, i) -entry of $(BB^\top)^{-1}$. See also Remark 3 (Originally, Remark 4.7 in [6]).

Considering the matrix $BB^\top$ instead of $B^\top B$, we have another recurrence relation as follows.

Theorem 11

Let B be an upper bidiagonal matrix defined in (1).

Let us introduce constants F_i for $i = 1, \dots, N-1$ defined as

$$F_i = \frac{e_i}{q_i}, \quad i = 1, \dots, N-1.$$

For $i = 1, \dots, N$ and $p = 1, 2, \dots$, let $\tilde{h}_i^{(p)}$ be constants which satisfy the following recurrence relation

$$\begin{aligned} \tilde{h}_N^{(1)} &= \frac{1}{q_N}, \\ \tilde{h}_i^{(1)} &= F_i \tilde{h}_{i+1}^{(1)} + \frac{1}{q_i}, \quad i = 1, \dots, N-1, \\ \tilde{h}_N^{(p)} &= 0, \quad p = 2, 3, \dots, \\ \tilde{h}_i^{(p)} &= F_i \left(\tilde{h}_{i+1}^{(p)} + p \tilde{h}_{i+1}^{(1)} \tilde{h}_{i+1}^{(p-1)} \right) \\ &\quad + \sum_{k=1}^{p-2} p C_k \tilde{h}_{i+1}^{(k)} \tilde{h}_i^{(p-k)}, \quad i = 1, \dots, N-1, \quad p = 2, 3, \dots \end{aligned}$$

For $i = 1, \dots, N$ and $p = 1, 2, \dots$, let $\tilde{H}_i^{(p)}$ be constants which satisfy the following recurrence relation

$$\begin{aligned} \tilde{H}_i^{(1)} &= \tilde{h}_i^{(1)}, \quad i = 1, \dots, N, \\ \tilde{H}_i^{(p)} &= \tilde{h}_i^{(p)} + \sum_{k=1}^{p-1} p C_k \tilde{h}_i^{(k)} \tilde{H}_i^{(p-k)}, \quad i = 1, \dots, N, \quad p = 2, 3, \dots \end{aligned}$$

The traces $\text{Tr}((BB^\top)^{-p})$ for $p = 1, 2, \dots$ are computed by

$$\text{Tr}((BB^\top)^{-p}) = \frac{1}{(p-1)!} \sum_{i=1}^N \tilde{H}_i^{(p)}. \quad (33)$$

Remark 12

The recurrence relation of $\tilde{h}_i^{(1)}$ ($i = 1, \dots, N$) in Theorem 11 is equivalent to the recurrence relation for the diagonal entries of $(B^\top B)^{-1}$ shown in Remark 2 (Originally, Remark 4.6 in [6]). Then, the constants $\tilde{h}_i^{(1)}$ and $\tilde{H}_i^{(1)}$ are the (i, i) -entry of $(B^\top B)^{-1}$.

The recurrence relations presented in Theorems 9 and 11 are subtraction-free.

3 A unified understanding of the two formulae in the preceding works and a new subtraction-free formula for the traces

In this section, we achieve a unified understanding of the two formulae in the preceding works [11] and [12] which are reviewed in the previous section. Firstly, we give an interpretation of some quantities, which is introduced in the formula in [11], in terms of matrix theory. Next, we show a relationship between some quantities in the formulae in [11] and [12]. Lastly, a new subtraction-free formula for the traces $J_M(B)$ ($M = 1, 2, \dots$) is naturally derived from this relationship.

Let us consider the quantities $\tilde{g}_i^{(M)}$ for $i = 1, \dots, N$ and $M = 1, 2, \dots$ given in Definition 6. We give an interpretation of the quantities $\tilde{g}_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ in terms of matrix theory. The following theorem holds.

Theorem 13

Let B be an upper bidiagonal matrix defined in (1). Let $W = (W_{i,j})$ denote the matrix $(BB^\top)^{-1}$. The quantities $\tilde{g}_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ which are given in Definition 6 are expressed as

$$\begin{aligned}\tilde{g}_i^{(2)} &= \sum_{j_1=1}^{i-1} W_{i,j_1} W_{j_1,i}, \\ \tilde{g}_i^{(3)} &= \sum_{j_1=1}^{i-1} \sum_{j_2=1}^{i-1} W_{i,j_1} W_{j_1,j_2} W_{j_2,i}, \\ \tilde{g}_i^{(M)} &= \sum_{j_1=1}^{i-1} \cdots \sum_{j_{M-1}=1}^{i-1} W_{i,j_1} W_{j_1,j_2} \cdots W_{j_{M-2},j_{M-1}} W_{j_{M-1},i}, \quad M = 4, 5, \dots\end{aligned}$$

Proof of this theorem is given in Appendix.

Next, we reveal a relationship between the quantities $\tilde{g}_i^{(M)}$ and $h_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$. As a preliminary, let us derive expressions for the traces $J_2(B)$ and $J_3(B)$ with $\tilde{g}_i^{(2)}$, $\tilde{g}_i^{(3)}$ and $W_{i,i}(= h_i^{(1)})$ ($i = 1, \dots, N$). The trace $J_2(B)$ is given as

$$J_2(B) = \sum_{i=1}^N \sum_{j=1}^N W_{i,j} W_{j,i}. \quad (34)$$

Let us classify the products $W_{i,j} W_{j,i}$ in (34) into three sets according to the subscripts i and j . Namely, i) $i > j$, ii) $i < j$ and iii) $i = j$. The summation

of the products which belong to the first set is

$$\sum_{i=1}^N \sum_{j=1}^{i-1} W_{i,j} W_{j,i} = \sum_{i=1}^N \tilde{g}_i^{(2)}$$

from Theorem 13. That of the second set is equal to this summation. That of the third set is

$$\sum_{i=1}^N (W_{i,i})^2 = \sum_{i=1}^N (h_i^{(1)})^2.$$

Then, the trace $J_2(B)$ is written as

$$J_2(B) = \sum_{i=1}^N \left(2\tilde{g}_i^{(2)} + (h_i^{(1)})^2 \right). \quad (35)$$

The trace $J_3(B)$ is given as

$$J_3(B) = \sum_{i=1}^N \sum_{j=1}^N \sum_{k=1}^N W_{i,j} W_{j,k} W_{k,i}. \quad (36)$$

The products $W_{i,j} W_{j,k} W_{k,i}$ in (36) are classified into seven sets according to the subscripts i , j and k . Namely, i) $i > j$ and $i > k$, ii) $j > i$ and $j > k$, iii) $k > i$ and $k > j$, iv) $i = j > k$, v) $i = k > j$, vi) $j = k > i$ and vii) $i = j = k$. The summation of the products which belong to the first set is

$$\sum_{i=1}^N \sum_{j=1}^{i-1} \sum_{k=1}^{i-1} W_{i,j} W_{j,k} W_{k,i} = \sum_{i=1}^N \tilde{g}_i^{(3)}$$

from Theorem 13. Those of the second and the third sets are equal to this summation. For the fourth set, the summation is

$$\sum_{i=1}^N \sum_{k=1}^{i-1} W_{i,i} W_{i,k} W_{k,i} = \sum_{i=1}^N h_i^{(1)} \tilde{g}_i^{(2)}$$

from Theorem 13. Those of the fifth and the sixth sets are equal to this summation. For the seventh set, the summation is

$$\sum_{i=1}^N (W_{i,i})^3 = \sum_{i=1}^N (h_i^{(1)})^3.$$

Thus, the trace $J_3(B)$ is expressed as

$$J_3(B) = \sum_{i=1}^N \left(3\tilde{g}_i^{(3)} + 3\tilde{g}_i^{(2)} h_i^{(1)} + (h_i^{(1)})^3 \right). \quad (37)$$

On the other hand, it is possible to express these traces with $h_i^{(1)}$, $h_i^{(2)}$ and $h_i^{(3)}$ ($i = 1, \dots, N$). From Theorem 9, we readily have

$$J_2(B) = \sum_{i=1}^N \left(h_i^{(2)} + \left(h_i^{(1)} \right)^2 \right), \quad (38)$$

$$J_3(B) = \frac{1}{2} \sum_{i=1}^N \left(h_i^{(3)} + 3h_i^{(2)}h_i^{(1)} + 2\left(h_i^{(1)} \right)^3 \right). \quad (39)$$

Comparison of (35) with (38) and of (37) with (39) suggests us to investigate whether it holds that

$$h_i^{(M)} = M! \cdot \tilde{g}_i^{(M)}, \quad i = 1, \dots, N, \quad M = 2, 3, \dots,$$

or not. The answer of the investigation is “true”. The following theorem holds.

Theorem 14

For $i = 1, \dots, N$ and $M = 2, 3, \dots$, the transform between a quantities $\tilde{g}_i^{(M)}$ and $h_i^{(M)}$ is given as

$$h_i^{(M)} = M! \cdot \tilde{g}_i^{(M)}. \quad (40)$$

Proof of this theorem is given in Appendix.

Now, we derive the new subtraction-free formula. Let us introduce the following transform

$$H_i^{(M)} = (M-1)! \cdot \tilde{G}_i^{(M)}, \quad i = 1, \dots, N, \quad M = 1, 2, \dots \quad (41)$$

Then, we readily have a new formula for the traces $J_M(B)$ ($M = 1, 2, \dots$) as follows by substituting (40) and (41) into the recurrence relation in Theorem 9 in Section 2.3.

Theorem 15

Let B be an $N \times N$ upper bidiagonal matrix defined as (1). Let $\check{B}_i$ for $i = 1, \dots, N$ be quantities defined by (14). Let $\tilde{F}_i$ for $i = 2, \dots, N$ be quantities defined by (16).

Let $\tilde{g}_i^{(1)}$ and $\tilde{G}_i^{(1)}$ for $i = 1, \dots, N$ be quantities which satisfy the recurrence relation

$$\tilde{g}_1^{(1)} = 0, \quad (42)$$

$$\tilde{G}_1^{(1)} = \check{B}_1, \quad (43)$$

$$\tilde{g}_i^{(1)} = \tilde{F}_i \tilde{G}_{i-1}^{(1)}, \quad i = 2, \dots, N, \quad (44)$$

$$\tilde{G}_i^{(1)} = \tilde{g}_i^{(1)} + \check{B}_i, \quad i = 2, \dots, N. \quad (45)$$

Let $\tilde{g}_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ be quantities which satisfy the following recurrence relation

$$\tilde{g}_1^{(M)} = 0, \quad (46)$$

$$\tilde{g}_i^{(M)} = \tilde{F}_i \tilde{g}_{i-1}^{(M)} + \tilde{G}_{i-1}^{(1)} \tilde{g}_i^{(M-1)} + \sum_{k=2}^{M-1} \tilde{g}_{i-1}^{(k)} \tilde{g}_i^{(M-k)}, \quad i = 2, \dots, N. \quad (47)$$

Let $\tilde{G}_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ be quantities which satisfy the following recurrence relation

$$\tilde{G}_i^{(M)} = M \tilde{g}_i^{(M)} + \tilde{G}_i^{(1)} \tilde{G}_i^{(M-1)} + \sum_{k=2}^{M-1} \tilde{g}_i^{(k)} \tilde{G}_i^{(M-k)}. \quad (48)$$

Then, the trace $J_M(B)$ for an arbitrary positive integer M is obtained from

$$J_M(B) = \sum_{i=1}^N \tilde{G}_i^{(M)}. \quad (49)$$

Proof of this theorem is given in Appendix.

As a comparison of the formulae in Theorems 9 and 15, we have the following remark.

Remark 16

In contrast to the formula in Theorem 9, the formula in Theorem 15 includes neither combination nor factorial in it. The quantities $\tilde{G}_i^{(M)}$ ($i = 1, \dots, N$) in Theorem 15 directly gives the trace $J_M(B)$ as their summation in contrast to the quantities $H_i^{(M)}$ ($i = 1, \dots, N$) in Theorem 9.

The relationship (41) for $M = 1$ and Remark 10 give the following remark.

Remark 17

The quantities $\tilde{G}_i^{(1)}$ ($i = 1, \dots, N$) are the (i, i) -entry of $(BB^\top)^{-1}$.

Let us consider the quantities $g_i^{(M)}$ for $i = 1, \dots, N$ and $M = 1, 2, \dots$ given in Definition 5 instead of $\tilde{g}_i^{(M)}$. Similarly to above, we have the following theorems and remarks.

Theorem 18

Let B be an upper bidiagonal matrix defined in (1). Let $V = (V_{i,j})$ denote the matrix $(B^\top B)^{-1}$. The quantities $g_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$

which are given in Definition 5 are expressed as

$$\begin{aligned}
g_i^{(2)} &= \sum_{j_1=i+1}^N V_{i,j_1} V_{j_1,i}, \\
g_i^{(3)} &= \sum_{j_1=i+1}^N \sum_{j_2=i+1}^N V_{i,j_1} V_{j_1,j_2} V_{j_2,i}, \\
g_i^{(M)} &= \sum_{j_1=i+1}^N \cdots \sum_{j_{M-1}=i+1}^N V_{i,j_1} V_{j_1,j_2} \cdots V_{j_{M-2},j_{M-1}} V_{j_{M-1},i}, \quad M = 4, 5, \dots
\end{aligned}$$

Theorem 19

For $i = 1, \dots, N$ and $M = 2, 3, \dots$, a transform between the quantities $g_i^{(M)}$ and $\tilde{h}_i^{(M)}$ is given as

$$\tilde{h}_i^{(M)} = M! \cdot g_i^{(M)}.$$

Theorem 20

Let B be an $N \times N$ upper bidiagonal matrix defined as (1). Let $\check{B}_i$ for $i = 1, \dots, N$ be quantities defined by (14). Let F_i for $i = 1, \dots, N - 1$ be quantities defined by (15).

Let $g_i^{(1)}$ and $G_i^{(1)}$ for $i = 1, \dots, N$ be quantities which satisfy the recurrence relation

$$\begin{aligned}
g_N^{(1)} &= 0, \\
G_N^{(1)} &= \check{B}_N, \\
g_i^{(1)} &= F_i G_{i+1}^{(1)}, & i = 1, \dots, N - 1, \\
G_i^{(1)} &= g_i^{(1)} + \check{B}_i, & i = 1, \dots, N - 1.
\end{aligned}$$

Let $g_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ be quantities which satisfy the following recurrence relation

$$\begin{aligned}
g_N^{(M)} &= 0, \\
g_i^{(M)} &= F_i g_{i+1}^{(M)} + G_{i+1}^{(1)} g_i^{(M-1)} + \sum_{k=2}^{M-1} g_{i+1}^{(k)} g_i^{(M-k)}, \quad i = 2, \dots, N.
\end{aligned}$$

Let $G_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ be quantities which satisfy the following recurrence relation

$$G_i^{(M)} = M g_i^{(M)} + G_i^{(1)} G_i^{(M-1)} + \sum_{k=2}^{M-1} g_i^{(k)} G_i^{(M-k)}.$$

Then, the trace $J_M(B)$ for an arbitrary positive integer M is obtained from

$$J_M(B) = \sum_{i=1}^N G_i^{(M)}.$$

Remark 21

In contrast to the formula in Theorem 11, the formula in Theorem 20 includes neither combination nor factorial in it. The quantities $G_i^{(M)}$ ($i = 1, \dots, N$) in Theorem 20 directly gives the trace $J_M(B)$ as their summation in contrast to the quantities $\tilde{H}_i^{(M)}$ ($i = 1, \dots, N$) in Theorem 11.

Remark 22

The quantities $G_i^{(1)}$ ($i = 1, \dots, N$) are the (i, i) -entry of $(B^\top B)^{-1}$.

4 Concluding Remarks

In this paper, we achieve a unified understanding of the formulae for the traces of the inverse powers of a positive definite symmetric tridiagonal matrix in the two preceding works. We show that the quantities $g_i^{(r)}$ and $\tilde{g}_i^{(r)}$ which are introduced in [11] and are shown in Section 2.2 can be interpreted in terms of matrix theory. The relationships which hold between some quantities in the two formulae in [11] and [12] are revealed. From these relationships, a new subtraction-free formula for the traces is obtained.

References

- [1] K. V. Fernando and B. N. Parlett, Accurate singular values and differential qd algorithms, *Numer. Math.*, **67** (1994) 191–229.
- [2] Y. P. Hong and C.-T. Pan, A lower bound for the smallest singular value, *Linear Algebra Appl.* **172** (1992) 27–32.
- [3] M. Iwasaki and Y. Nakamura, Accurate computation of singular values in terms of shifted integrable schemes, *Japan J. Indust. Appl. Math.*, **23** (2006) 239–259.
- [4] C. R. Johnson, A Gersgorin-type lower bound for the smallest singular value, *Linear Algebra Appl.* **112** (1989) 1–7.
- [5] C. R. Johnson and T. Szulc, Further lower bounds for the smallest singular value, *Linear Algebra Appl.* **272** (1998) 169–179.

- [6] K. Kimura, T. Yamashita and Y. Nakamura, Conserved quantities of the discrete finite Toda equation and lower bounds of the minimal singular value of upper bidiagonal matrices, *J. Phys. A: Math. Theor.*, **44** (2011) 285207 (12pp.).
- [7] L. Li, Lower bounds for the smallest singular value, *Computers and Mathematics with Applications*, **41** (2001) 483–487.
- [8] U. von Matt, The orthogonal qd-algorithm, *SIAM J. Sci. Comput.*, **18** (1997) 1163–1186.
- [9] O. Rojo, Further bounds for the smallest singular value and the spectral condition number, *Computers and Mathematics with Applications*, **38** (1999) 215–228.
- [10] H. Rutishauser, *Lectures on Numerical Mathematics*, Birkhäuser Boston, Cambridge, MA, 1990.
- [11] T. Yamashita, K. Kimura and Y. Nakamura, Subtraction-free recurrence relations for lower bounds of the minimal singular value of an upper bidiagonal matrix, *Journal of Math-for-Industry*, **4** (2012) 55–71.
- [12] T. Yamashita, K. Kimura and Y. Yamamoto, A new subtraction-free formula for lower bounds of the minimal singular value of an upper bidiagonal matrix, *Numer. Algor.*, accepted, DOI: 10.1007/s11075-014-9931-z.
- [13] Y.-S. Yu and D.-H. Gu, A note on a lower bound for the smallest singular value, *Linear Algebra Appl.*, **253** (1997) 25–38.

Appendix

In this appendix, we give proof of Theorems 13, 14 and 15. For proof of Theorems 13 and 14, we give the following remark.

Remark 23

For $i = 2, \dots, N$, the quantities $\tilde{g}_i^{(1)}$ can be written as

$$\tilde{g}_i^{(1)} = \tilde{F}_i W_{i-1,i-1} \tag{50}$$

from its definition. For $i = 2, \dots, N$, we can readily verify that it holds $\tilde{g}_{i-1}^{(1)} + \tilde{B}_{i-1} = W_{i-1,i-1}$ from Definition 6 and Remark 7. Then, the recurrence

relation (18) in Definition 6 for $i = 2, \dots, N$ and $M = 2, 3, \dots$ can be rewritten as

$$\tilde{g}_i^{(M)} = \tilde{F}_i \tilde{g}_{i-1}^{(M)} + W_{i-1, i-1} \tilde{g}_i^{(M-1)} + \sum_{k=2}^{M-1} \tilde{g}_{i-1}^{(k)} \tilde{g}_i^{(M-k)}. \quad (51)$$

Before showing proof of Theorem 13, we give a lemma for this proof.

Lemma 24

If $i > j$ and $i > k$, then it holds

$$W_{i,j} W_{k,i} = \tilde{F}_i W_{i-1,j} W_{k,i-1}, \quad (52)$$

$$W_{j,i-1} W_{k,i} = W_{j,i} W_{k,i-1}. \quad (53)$$

proof

Let $S = (S_{i,j})$ denote the inverse of B . Then, the matrix W is expressed as $W = S^\top S$ from the definition $W = (BB^\top)^{-1}$. This matrix S is an upper triangle matrix. The relationship

$$S_{i,j} = -\sqrt{\tilde{F}_j} \cdot S_{i,j-1}, \quad 1 \leq i < j \leq N$$

holds among entries of S (See [6].). This relationship gives relationships among entries of W . When $i > j$, since it holds

$$\begin{aligned} W_{i,j} &= \sum_{k=1}^N (S^\top)_{i,k} S_{k,j} = \sum_{k=1}^N S_{k,i} S_{k,j} = \sum_{k=1}^j S_{k,i} S_{k,j} \\ &= \sum_{k=1}^j \left(-\sqrt{\tilde{F}_i} \cdot S_{k,i-1} \right) S_{k,j} = -\sqrt{\tilde{F}_i} \sum_{k=1}^N (S^\top)_{i-1,k} S_{k,j} \\ &= -\sqrt{\tilde{F}_i} \cdot W_{i-1,j}, \end{aligned}$$

we have

$$W_{i,j} = -\sqrt{\tilde{F}_i} \cdot W_{i-1,j}.$$

Since W is a symmetric matrix, it holds

$$W_{j,i} = -\sqrt{\tilde{F}_i} \cdot W_{j,i-1}$$

when $i > j$. Then, when $i > j$ and $i > k$, it holds (52) and (53). $\square$

Proof of Theorem 13

Let us introduce quantities $u_i^{(M)}$ for $i = 1, \dots, N$ and $M = 1, 2, \dots$ as follows. The quantities $u_i^{(1)}$ ($i = 1, \dots, N$) are defined as

$$u_i^{(1)} = \begin{cases} 0, & i = 1, \\ \tilde{F}_i W_{i-1, i-1}, & i = 2, \dots, N. \end{cases}$$

The quantities $u_i^{(M)}$ for $i = 1, \dots, N$ and $M = 2, 3, \dots$ are defined as

$$\begin{aligned} u_i^{(2)} &= \sum_{j_1=1}^{i-1} W_{i, j_1} W_{j_1, i}, \\ u_i^{(3)} &= \sum_{j_1=1}^{i-1} \sum_{j_2=1}^{i-1} W_{i, j_1} W_{j_1, j_2} W_{j_2, i}, \\ u_i^{(r)} &= \sum_{j_1=1}^{i-1} \cdots \sum_{j_{r-1}=1}^{i-1} W_{i, j_1} W_{j_1, j_2} \cdots W_{j_{r-2}, j_{r-1}} W_{j_{r-1}, i}, \quad r = 4, 5, \dots \end{aligned}$$

For $M = 1, 2, \dots$, the quantities $u_1^{(M)}$ and $\tilde{g}_1^{(M)}$ coincide since they are zero. From the definitions, the quantities $u_i^{(1)}$ and $\tilde{g}_i^{(1)}$ coincide for $i = 2, \dots, N$. Then, when we show that a recurrence relation which the quantities $u_i^{(M)}$ for $i = 2, \dots, N$ and $M = 2, 3, \dots$ satisfy has the same form as (51) in Remark 23, proof completes. We consider the cases where $M = 2$. It holds $\tilde{g}_1^{(2)} = u_1^{(2)}$ as mentioned above. The quantities $u_i^{(2)}$ for $i = 2, \dots, N$ satisfy

$$\begin{aligned} u_i^{(2)} &= \sum_{j=1}^{i-1} W_{i, j} W_{j, i} = \sum_{j=1}^{i-1} \tilde{F}_i W_{i-1, j} W_{j, i-1} \\ &= \tilde{F}_i \sum_{j=1}^{i-2} W_{i-1, j} W_{j, i-1} + \tilde{F}_i W_{i-1, i-1} W_{i-1, i-1} \\ &= \tilde{F}_i u_{i-1}^{(2)} + W_{i-1, i-1} u_i^{(1)} \end{aligned} \tag{54}$$

from (52) in Lemma 24 and the definition of $u_i^{(1)}$. Then, the recurrence relations (51) for $M = 2$ in Remark 23 and (54) have the same form. Thus, $\tilde{g}_i^{(2)}$ and $u_i^{(2)}$ for $i = 1, \dots, N$ coincide.

Next, we consider the cases where $i = 2$. As mentioned above, it holds $\tilde{g}_2^{(M)} = u_2^{(M)}$ for $M = 1$ and 2. Then let us consider cases where $M = 3, 4, \dots$. It holds $\tilde{g}_2^{(M)} = W_{1,1} \tilde{g}_2^{(M-1)}$ from (51) in Remark 23 since $\tilde{g}_1^{(l)} = 0$ ($l = 2, \dots, M$). It holds $u_2^{(M)} = W_{2,1} (W_{1,1})^{M-2} W_{1,2} = W_{1,1} u_2^{(M-1)}$ from the definition. Then, for $M = 1, 2, \dots$, the quantities $\tilde{g}_2^{(M)}$ and $u_2^{(M)}$ coincide.

Lastly, let us consider cases where M, N and i are $M \geq 3, N \geq 3$ and $i = 3, \dots, N$, respectively. Let us divide the products $W_{i,j_1} \cdots W_{j_{M-1},i}$ which appear in the definition of $u_i^{(M)}$ into M sets $S_0, S_1, \dots, S_{M-1}$. The way of division is as follows. If the suffixes $j_1, \dots, j_{M-1}$ are not $i-1$, then the product belongs to the set S_0 . If the suffix j_{M-1} is $i-1$, then the product belongs to the set S_{M-1} . If the suffix j_k is $j_k = i-1$ for some k ($1 \leq k \leq M-2$) and the suffix(es) j_l for $l = k+1, \dots, M-1$ is/are $j_l < i-1$, then the product belongs to the set S_k . Let the summation of the products which belong to the set S_k ($k = 0, 1, \dots, M-1$) be denoted by $\alpha_i^{(M,k)}$. Then, the summation $u_i^{(M)}$ is given as

$$u_i^{(M)} = \sum_{k=0}^{M-1} \alpha_i^{(M,k)}. \quad (55)$$

The summation $\alpha_i^{(M,0)}$ is written as

$$\begin{aligned} \alpha_i^{(3,0)} &= \sum_{j_1=1}^{i-2} \sum_{j_2=1}^{i-2} W_{i,j_1} W_{j_1,j_2} W_{j_2,i}, \\ \alpha_i^{(r,0)} &= \sum_{j_1=1}^{i-2} \cdots \sum_{j_{r-1}=1}^{i-2} W_{i,j_1} W_{j_1,j_2} \cdots W_{j_{r-2},j_{r-1}} W_{j_{r-1},i}, \quad r = 4, 5, \dots \end{aligned}$$

Since it holds $W_{i,j_1} W_{j_{M-1},i} = \tilde{F}_i W_{i-1,j_1} W_{j_{M-1},i-1}$ from (52) in Lemma 24, we have

$$\alpha_i^{(M,0)} = \tilde{F}_i u_{i-1}^{(M)}. \quad (56)$$

The summation $\alpha_i^{(M,M-1)}$ is written as

$$\begin{aligned} \alpha_i^{(3,2)} &= \sum_{j_1=1}^{i-1} W_{i,j_1} W_{j_1,i-1} W_{i-1,i}, \\ \alpha_i^{(4,3)} &= \sum_{j_1=1}^{i-1} \sum_{j_2=1}^{i-1} W_{i,j_1} W_{j_1,j_2} W_{j_2,i-1} W_{i-1,i}, \\ \alpha_i^{(r,r-1)} &= \sum_{j_1=1}^{i-1} \cdots \sum_{j_{r-2}=1}^{i-1} W_{i,j_1} W_{j_1,j_2} \cdots W_{j_{r-2},i-1} W_{i-1,i}, \quad r = 5, 6, \dots \end{aligned}$$

It holds $W_{j_{M-2},i-1} W_{i-1,i} = W_{j_{M-2},i} W_{i-1,i-1}$ from (53) in Lemma 24. Then, we have

$$\alpha_i^{(M,M-1)} = W_{i-1,i-1} u_i^{(M-1)}. \quad (57)$$

The summation $\alpha_i^{(M,1)}$ is written as

$$\begin{aligned}\alpha_i^{(3,1)} &= \sum_{j_2=1}^{i-2} W_{i,i-1} W_{i-1,j_2} W_{j_2,i}, \\ \alpha_i^{(4,1)} &= \sum_{j_2=1}^{i-2} \sum_{j_3=1}^{i-2} W_{i,i-1} W_{i-1,j_2} W_{j_2,j_3} W_{j_3,i}, \\ \alpha_i^{(r,1)} &= \sum_{j_2=1}^{i-2} \cdots \sum_{j_{r-1}=1}^{i-2} W_{i,i-1} W_{i-1,j_2} \cdots W_{j_{r-2},j_{r-1}} W_{j_{r-1},i}, \quad r = 5, 6, \dots\end{aligned}$$

Since it holds $W_{i,i-1} W_{j_{M-1},i} = \tilde{F}_i W_{i-1,i-1} W_{j_{M-1},i-1} = u_i^{(1)} W_{j_{M-1},i-1}$ from (52) in Lemma 24 and the definition of $u_i^{(1)}$, we have

$$\alpha_i^{(M,1)} = u_i^{(1)} u_{i-1}^{(M-1)}. \quad (58)$$

In the case of $M \geq 4$, the summation $\alpha_i^{(M,k)}$ ($k = 2, \dots, M-2$) is given as

$$\alpha_i^{(M,k)} = \sum_{j_1=1}^{i-1} \cdots \sum_{j_{k-1}=1}^{i-1} \sum_{j_{k+1}=1}^{i-2} \cdots \sum_{j_{M-1}=1}^{i-2} W_{i,j_1} \cdots W_{j_{k-1},i-1} W_{i-1,j_{k+1}} \cdots W_{j_{M-1},i}.$$

Since it holds $W_{j_{k-1},i-1} W_{j_{M-1},i} = W_{j_{k-1},i} W_{j_{M-1},i-1}$ from (53) in Lemma 24, we have

$$\alpha_i^{(M,k)} = u_i^{(k)} u_{i-1}^{(M-k)}. \quad (59)$$

From (55), (56), (57), (58) and (59), we obtain

$$\begin{aligned}u_i^{(M)} &= \sum_{k=0}^{M-1} \alpha_i^{(M,k)} = \tilde{F}_i u_{i-1}^{(M)} + W_{i-1,i-1} u_i^{(M-1)} + \sum_{k=1}^{M-2} u_i^{(k)} u_{i-1}^{(M-k)} \\ &= \tilde{F}_i u_{i-1}^{(M)} + W_{i-1,i-1} u_i^{(M-1)} + \sum_{k=2}^{M-1} u_{i-1}^{(k)} u_i^{(M-k)}.\end{aligned} \quad (60)$$

Then, the recurrence relations (51) in Remark 23 and (60) have the same form. Thus, $\tilde{g}_i^{(M)}$ and $u_i^{(M)}$ for $i = 3, \dots, N$ and $M = 3, 4, \dots$ coincide. $\square$

Proof of Theorem 14

Let us introduce quantities $\zeta_i^{(M)}$ for $i = 1, \dots, N$ and $M = 1, 2, \dots$ defined as

$$\zeta_1^{(1)} = 0, \quad (61)$$

$$\zeta_i^{(1)} = \tilde{F}_i h_{i-1}^{(1)}, \quad i = 2, \dots, N, \quad (62)$$

$$\zeta_i^{(M)} = \frac{1}{M!} \cdot h_i^{(M)}, \quad i = 1, \dots, N, \quad M = 2, 3, \dots \quad (63)$$

When we show that a recurrence relation which the quantities $\zeta_i^{(M)}$ satisfy coincides with that of the quantities $\tilde{g}_i^{(M)}$ in Definition 6, proof completes.

It is obvious that $\zeta_1^{(M)} = \tilde{g}_1^{(M)}$ for $M = 1, 2, \dots$ since they are zero from their definitions.

Hereafter, let i be $i = 2, \dots, N$ in this proof.

The quantities $h_{i-1}^{(1)}$ coincide with the diagonal entry $W_{i-1,i-1}$ of W from Remark 10. Then, it holds

$$\zeta_i^{(1)} = \tilde{g}_i^{(1)}$$

from the definition of $\tilde{g}_i^{(1)}$.

Let us consider the case where $M = 2$. From (63), the recurrence relation for $h_i^{(2)}$ in Theorem 9 is rewritten as

$$2\zeta_i^{(2)} = \tilde{F}_i \left(2\zeta_{i-1}^{(2)} + 2 \left(h_{i-1}^{(1)} \right)^2 \right).$$

From (62) and the coincidence $h_{i-1}^{(1)} = W_{i-1,i-1}$, we have

$$\zeta_i^{(2)} = \tilde{F}_i \zeta_{i-1}^{(2)} + W_{i-1,i-1} \zeta_i^{(1)}.$$

This recurrence relation has the same form with the recurrence relation (51) for $M = 2$ in Remark 23.

Let us consider the case where $M = 3, 4, \dots$. The recurrence relation (31) for $h_i^{(M)}$ in Theorem 9 is rewritten as

$$h_i^{(M)} = \tilde{F}_i h_{i-1}^{(M)} + M \tilde{F}_i h_{i-1}^{(1)} h_{i-1}^{(M-1)} + M h_{i-1}^{(1)} h_i^{(M-1)} + \sum_{k=2}^{M-2} {}_M C_k h_{i-1}^{(k)} h_i^{(M-k)}.$$

On the second and the third terms in the right-hand-side, it holds

$$\begin{aligned} & M \tilde{F}_i h_{i-1}^{(1)} h_{i-1}^{(M-1)} + M h_{i-1}^{(1)} h_i^{(M-1)} \\ &= M \zeta_i^{(1)} \cdot (M-1)! \cdot \zeta_{i-1}^{(M-1)} + M W_{i-1,i-1} \cdot (M-1)! \cdot \zeta_i^{(M-1)} \\ &= M! \cdot \left(W_{i-1,i-1} \zeta_i^{(M-1)} + \zeta_i^{(1)} \zeta_{i-1}^{(M-1)} \right). \end{aligned}$$

Then, we have

$$\begin{aligned} M! \cdot \zeta_i^{(M)} &= \tilde{F}_i \cdot M! \cdot \zeta_{i-1}^{(M)} + M! \cdot \left(W_{i-1,i-1} \zeta_i^{(M-1)} + \zeta_i^{(1)} \zeta_{i-1}^{(M-1)} \right) \\ &\quad + \sum_{k=2}^{M-2} {}_M C_k \cdot k! \cdot \zeta_{i-1}^{(k)} \cdot (M-k)! \cdot \zeta_i^{(M-k)} \end{aligned}$$

from the definition (63). Since it holds ${}_M C_k \cdot k! \cdot (M-k)! = M!$, we immediately have

$$\zeta_i^{(M)} = \tilde{F}_i \zeta_{i-1}^{(M)} + W_{i-1,i-1} \zeta_i^{(M-1)} + \sum_{k=2}^{M-1} \zeta_{i-1}^{(k)} \zeta_i^{(M-k)}.$$

This recurrence relation has the same form as (51) for $M = 3, 4, \dots$ in Remark 23. $\square$

Proof of Theorem 15

From Remarks 10 and 17, it holds

$$\tilde{G}_i^{(1)} = H_i^{(1)} = h_i^{(1)}, \quad i = 1, \dots, N. \quad (64)$$

Substituting $h_i^{(1)} = \tilde{G}_i^{(1)}$ ($i = 1, \dots, N$) into (28) and (29) in Theorem 9 and defining $\tilde{g}_i^{(1)}$ ($i = 1, \dots, N$) as (42) and (44), we readily have (43) and (45).

Next, we substitute (40) in Theorem 14 into (30) and (31) in Theorem 9. We readily have (46). And we have

$$\begin{aligned} M! \cdot \tilde{g}_i^{(M)} &= \tilde{F}_i \left(M! \cdot \tilde{g}_{i-1}^{(M)} + M \cdot h_{i-1}^{(1)} \cdot (M-1)! \cdot \tilde{g}_{i-1}^{(M-1)} \right) \\ &\quad + M \cdot h_{i-1}^{(1)} \cdot (M-1)! \cdot \tilde{g}_i^{(M-1)} \\ &\quad + \sum_{k=2}^{M-2} M C_k \cdot k! \cdot \tilde{g}_{i-1}^{(k)} \cdot (M-k)! \cdot \tilde{g}_i^{(M-k)} \end{aligned}$$

for $i = 2, \dots, N$ and $M = 2, 3, \dots$. Then, it holds

$$\tilde{g}_i^{(M)} = \tilde{F}_i \tilde{g}_{i-1}^{(M)} + \tilde{F}_i h_{i-1}^{(1)} \tilde{g}_{i-1}^{(M-1)} + h_{i-1}^{(1)} \tilde{g}_i^{(M-1)} + \sum_{k=2}^{M-2} \tilde{g}_{i-1}^{(k)} \tilde{g}_i^{(M-k)}.$$

The second term of the right-hand-side is equal to $\tilde{g}_i^{(1)} \tilde{g}_{i-1}^{(M-1)}$ since it holds $\tilde{F}_i h_{i-1}^{(1)} = \tilde{F}_i \tilde{G}_{i-1}^{(1)} = \tilde{g}_i^{(1)}$ from (44) and (64). From (64), the third term of the right-hand-side is equal to $\tilde{G}_{i-1}^{(1)} \tilde{g}_i^{(M-1)}$. Then, we have (47).

Next, we substitute (40) in Theorem 14 and (41) into (32) in Theorem 9. We have

$$\begin{aligned} (M-1)! \cdot \tilde{G}_i^{(M)} &= M! \cdot \tilde{g}_i^{(M)} + (M-1) h_i^{(1)} \cdot (M-2)! \cdot \tilde{G}_i^{(M-1)} \\ &\quad + \sum_{k=2}^{M-1} M_{-1} C_k \cdot k! \cdot \tilde{g}_i^{(k)} \cdot (M-1-k)! \cdot \tilde{G}_i^{(M-k)} \end{aligned}$$

for $i = 1, \dots, N$ and $M = 2, 3, \dots$. Then, it holds

$$\tilde{G}_i^{(M)} = M \tilde{g}_i^{(M)} + h_i^{(1)} \tilde{G}_i^{(M-1)} + \sum_{k=2}^{M-1} \tilde{g}_i^{(k)} \tilde{G}_i^{(M-k)}$$

for $i = 1, \dots, N$ and $M = 2, 3, \dots$. Since the second term of the right-hand-side is equal to $\tilde{G}_i^{(1)} \tilde{G}_i^{(M-1)}$ from (64), we obtain (48).

Lastly, substituting (41) into (33) in Theorem 9, we readily have (49).

$\square$